

1. Suppose the solutions of a matrix equation $A\mathbf{x} = \mathbf{b}$ are written in the form $\mathbf{p} + \mathbf{v}_h$, where \mathbf{p} is a particular solution of $A\mathbf{x} = \mathbf{b}$ and \mathbf{v}_h gives all solutions of the corresponding homogeneous equation $A\mathbf{x} = \mathbf{0}$.

$$\text{Suppose } \mathbf{b} = \begin{bmatrix} 2 \\ -13 \\ 0 \\ 2009 \end{bmatrix}, \mathbf{p} = \begin{bmatrix} 11 \\ 0 \\ -2 \\ 5 \\ 0 \end{bmatrix} \text{ and } \mathbf{v}_h = x_2 \begin{bmatrix} -9 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 7 \\ 0 \\ -2 \\ 0 \\ 1 \end{bmatrix}, \text{ where } x_2 \text{ and } x_5 \text{ are free.}$$

1a. How many *rows* does A have?

How many *columns*?

1b. Label the columns of A as $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_p$. Is the set $S = \{\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_p\}$ *linearly independent*? Explain in terms of the definition of linear independence.

1c. Write \mathbf{c}_3 as a linear combination of the other columns of A , or explain why this cannot be done.

1d. Write \mathbf{c}_4 as a linear combination of the other columns of A , or explain why this cannot be done.

1e. Show how to express \mathbf{b} as a linear combination of all p columns of A in such a way that none of the weights involved are 0.

2. Suppose $T : \mathbb{R}^a \rightarrow \mathbb{R}^z$ is a transformation. Give the definitions of each of the following:

2a. T is a *linear* transformation.

2b. T is onto \mathbb{R}^z .

2c. Suppose $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ is defined by $T \left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \right) = \begin{bmatrix} x_2x_3 + 2x_1 \\ 0 \\ x_1 + x_2 + x_3 \\ 2x_2 + 7 \end{bmatrix}$. Show by example that T is not a linear transformation and that it actually fails both parts of the definition in (2a).

3. Suppose $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ is the linear transformation whose standard matrix is $A = \begin{bmatrix} 5 & 11 & 23 \\ 5 & 14 & 17 \\ 2 & 6 & 6 \\ 3 & 8 & 11 \end{bmatrix}$.

3a. Find $T \left(\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right)$.

3b. Let $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}$ be a vector in \mathbb{R}^4 . What, if any, conditions on b_1, \dots, b_4 must be met to guarantee

\mathbf{b} is in the range of T ? Show any matrices you use in answering this question.

3c. Suppose $\mathbf{d} = \begin{bmatrix} 3 \\ 12 \\ d_3 \\ d_4 \end{bmatrix}$. Use the conditions in (3b) to find all values of d_3 and d_4 for which \mathbf{d} is in

the range of T . (Note you will be setting up a little linear system, and you should use our linear algebra techniques to solve it).

3d. Is T onto \mathbb{R}^4 ? Explain your answer.

3e. Is T one-to-one? Explain your answer.

4. Let $F = \begin{bmatrix} 2 & -3 & w \\ 1 & 5 & -2 \end{bmatrix}$ and $G = \begin{bmatrix} 8 & x & -6 \\ 7 & -2 & 1 \\ 4 & y & -5 \end{bmatrix}$, and $P = \begin{bmatrix} 7 & 23 & z \\ q & -12 & 9 \end{bmatrix}$; suppose $FG = P$.

Show all your work in the following:

4a. Find q .

4b. Find w .

4c. Find z .

4d. Find x and y . Use linear algebra techniques to solve any system this problem requires.

5. Suppose that A is a 2×2 matrix and the following row operations convert A into I_2 : First, rows 1 and 2 of A are swapped. Then 4 copies of row 1 are subtracted from row 2. Finally, row 2 is multiplied by 5.

5a. What three elementary matrices E_1 , E_2 , and E_3 represent these three row operations, respectively?

5b. Use the elementary matrices to find A^{-1} .

5c. Find A .

5d. What is the determinant of A ?

5e. Is A singular or nonsingular?